Organic Corn Performance Test

A. Sundermeier, Extension Agent, Wood County

D.H. Stinner, Research Scientist, Dept. of Entomology

R.J. Minyo, Research Associate, Dept. of Horticulture & Crop Science

A.B. Geyer, Research Associate, Dept. of Horticulture & Crop Science

P.R. Thomison, Associate Professor, Dept. of Horticulture & Crop Science

Objective

Interest in growing corn in organic cropping systems has been on the rise in recent years. An evaluation of corn hybrids marketed for organic producers was conducted at the Hirzel Sustainable Systems site, a certified organic farm located near Bowling Green in Wood county. The data from this performance test is intended to help organic corn producers select hybrids to grow on their farms.

Background

Location: County: Nearest Town: Soil Type: Tillago:	Hirzel Sustainable Systems Site Wood Bowling Green Hoytville Clay Loam	Previous Crop: Planting Date: Planting Rate: Row Width: Harvest Date:	Alfalfa 5/28/04 23,000 seeds/A 30-inch 11/22/04
Tillage:	Conventional	Harvest Date:	11/22/04

Method

Plots were planted in a randomized complete block design with four reps. Four row plots were planted, with the center two rows harvested. Plots were 25 feet in length. No fertilizers or pesticides were applied to the site. Weed control was provided by mechanical cultivation. Notes and measurements were taken on the following: yield (bushels per acre at 15.5% moisture), harvest moisture, stalk lodging, final stand, percent emergence, test weight and silk day (the Julian day of the year in which 50% of the plants were silked).

Results

Performance data for the 2004 Organic Corn Performance Test.

Brand	Hybrid	Yield	Harv Moist	Stk Ldg	Final Stand	Emerg	Test Wt	Silk
		-Bu/A-	%-		Plants/A	%	lbs/bu	day
Breyleys	4920	93.0	21.7	16	20200	89	53.7	218
Breyleys	4930	105.3	21.4	32	24500	90	54.3	220
Breyleys	5720	104.8	22.7	30	22300	85	53.6	220
Breyleys	5740	102.3	21.7	40	23000	96	53.8	221
Doebler	75X2	101.5	28.5	18	26400	92	49.0	230
Doebler	N509	115.6	21.9	9	19100	93	54.8	219
Doebler	N640	103.5	24.1	15	24700	84	53.7	223
Great Harvest	41X2	116.1	21.8	15	22800	83	54.8	220
Great Harvest	43V6	104.3	21.6	43	23100	80	51.2	219
Great Harvest	44X2	118.0	23.4	13	24600	86	52.9	220
Great Harvest	56V6	118.3	22.4	18	21900	88	52.1	218
Great Harvest	60V6	110.5	25.3	48	26500	92	49.5	222
Merit Seeds	335	87.7	19.3	55	26100	88	56.2	218
Merit Seeds	535	114.2	22.6	23	20900	75	53.6	219
Merit Seeds	635	121.8	24.4	30	24700	83	53.4	223
NC+	081A2ORG	74.6	19.6	26	26300	90	58.6	213
NC+	0850ORG	86.2	19.7	22	26200	89	56.8	217
NC+	100A2ORG	99.6	20.4	38	27200	92	55.2	221
NC+	102A2ORG	100.7	22.0	14	25000	84	54.7	221
NC+	112E1ORG	109.6	25.8	19	24700	85	51.5	222
NC+	17A21ORG	90.4	19.6	15	25600	83	56.9	217
NC+	30A12ORG	91.4	19.8	35	28500	95	56.7	217
NC+	31F10ORG	88.6	19.8	56	27700	92	55.6	218
NC+	3448ORG	123.6	22.5	21	25800	89	53.8	217
NC+	34C17ORG	103.0	18.8	19	25900	93	55.6	217
NC+	40M21ORG	105.7	19.3	24	25900	86	56.0	217
NC+	42A32ORG	118.9	19.4	32	24300	86	54.5	217
NC+	4771ORG	108.0	21.5	40	24400	85	52.2	224
NC+	4822ORG	114.4	24.7	23	22200	76	50.3	223
NC+	48F37ORG	115.5	21.6	22	24900	83	53.6	218
NC+	49M37ORG	122.4	21.3	14	26000	88	53.6	219
NC+	50K32ORG	116.8	22.0	20	25900	93	54.3	218
NC+	51A30ORG	114.7	20.5	35	26000	92	54.4	220
NC+	54M52ORG	117.6	22.3	26	26800	96	53.4	220
NC+	57F00ORG	109.2	24.6	16	22600	80	52.6	223
NC+	57K36ORG	113.9	23.9	46	25100	91	51.1	223
NC+	58T36ORG	122.6	22.7	44	26200	91	51.3	222
NC+	68F32ORG	114.5	24.3	31	26200	86	53.8	221
NC+	68T57ORG	118.5	24.7	52	26000	92	50.2	225
NC+	72H54ORG	116.1	25.5	19	24800	93	48.8	223
		400.0						
HIGH		123.6	28.5	56	28500	96	58.6	230
AVERAGE		107.8	22.2	28	24800	88	53.5	220
		/4.6	18.8	9	19100	/5	48.8	213
LSD (0.05)		14.8	0.8	23	3000	12	0.9	2
CV		9.6	2.6	58	8	9	1.2	1

Summary

Growing conditions were generally favorable for corn production at the site, however, excessive rainfall in May delayed planting. Grain yields ranged from 74.6 to 123.6 bushels/acre; stalk lodging from 9-56%; and grain moisture at harvest from 18.8-28.5%. The relative maturity of the hybrids entered ranged from 81 to 116 days.

Acknowledgements/Seed Sources

For further information on specific hybrids, please contact the participating companies:						
Brand	Address	Hybrid	Relative Maturity*			
Doeblers	8999 E. Lincoln Way	75X2	116			
	Orrville, OH 44667	N509	101			
	330-683-3220	N460	111			
Broyleys	7007 New London Eastern Rd	1020	03			
Dieyley3	Spencer OH 44275	4920	100			
	330-667-2472	5720	105			
	555 567 2472	5740	108			
		0140	100			
	eese E i ezeth ei	40/0	4.04			
Great Harvest	6803 East 276" St	41 X2	101			
	Atlanta, IN 46031	4376	103			
	317-984-6685	44X2	104			
		56V6	105			
		60V6	110			
Merit Seeds	PO Box 205	335	91			
	Berlin, OH 44610	535	105			
	800-553-4713	635	108			
NC L Organica	DO Boy 4720	00140000	04			
NC+ Organics	2020 N 56 th St	001A20KG	01			
	Lincoln NE 68504	10042080	100			
	200 270 7000	100A20KG	100			
	000-279-7999	102A2ORG	102			
		174210RG	112			
		204120RG	04			
		30A120RG	90			
		31F100RG	91			
		3440URG	106			
		340170RG	93			
		401VIZTURG	90			
		42A32UKG	97			
		74410RG	110			
		4022URG	112			
		40F3/UKG	102			
		49101370RG	102			
			103			
		5TA3UUKG	103			
		541VI52UKG	105			
		5/FUUUKG	107			
		5/K30UKG	107			
		58136UKG	108			
		OOT SZUKG	112			
		6815/UKG	113			
		72H54ORG	114			

* Relative maturity ratings provided by companies.

For additional information, contact:

Peter Thomison The Ohio State University Horticulture and Crop Science 2021 Coffey Rd Columbus, OH 43210 614-292-2373 thomison.1@osu.edu