

## Soybean Seeding Rate

Eric Richer, Ohio State University Extension Educator, Fulton County

### Objective

To determine the effects of seeding rate on soybean yield and profitability.

### Background

Crop Year:	2015	Tillage:	Minimum
Location:	Archbold, OH	Soil Test (avg):	pH 6.7, P 139 ppm*
County:	Fulton		K 303 ppm, CEC 15.9
Soil Type:	Fulton & Latty clay		O.M. 2.8%
Drainage:	Systematic, 50' laterals	Planting Date:	May 8, 2014
Previous Crop:	Corn	Fertility:	Applied in corn year with VRT
Planting Date:	May 22, 2015	Harvest Date:	October 21, 2015
Herbicide:		Rainfall:	25.6" (Apr-Sept)
	3.5 oz Envive, 6 oz Glory, 1 pt 2,4d (Preplant)		*Mehlich-3 Extractant
	22 oz RoundUp PowerMax (Post Emerge)		

### Methods

Five treatments of different soybean populations were replicated four times in a randomized complete block design. Treatments were planted with a 40 foot John Deere 1790 air seeder in 15" rows. All treatments received the same tillage and herbicide applications. Seed used was Rupp 7332. Plot centers were harvested with a commercial combine equipped with a 35 foot grain header. Yields and moistures were obtained by using a calibrated GreenStar 2630 monitor. Yields were calculated at 13% moisture content. Precipitation data were obtained from the National Weather Service Wauseon WTP Station.

### Results

Seeding rate (x 1,000)	Yield Bu/acre	Gross Revenue per acre	Seed Cost per acre	Net Revenue per acre
97	46.1 cd	\$414.90	\$41.71	\$373.19
121	48.5 bc	\$436.50	\$52.03	\$384.47
145	51.7 ab	\$465.30	\$62.35	\$402.95
170	51.1 b	\$459.90	\$73.10	\$386.8
191	54.2 a	\$487.89	\$82.13	\$405.67
LSD (p<.05)	2.98 (cv 3.84)			

**Economics:** Gross income= yield x \$9.00/bu; Seed cost= \$0.43 per 1,000 seeds x seeding rate;  
Net revenue= Gross revenue – seed cost.



**Discussion:**

There was no statistically significant difference in yield (or economics) between the 145,000 and 191,000 seeds per acre treatments, however, those seeding rates yielded significantly better than any of the other three rates. Further data in the form of multi-year replications will add to the validity of these results.

**Acknowledgement**

The author expresses appreciation to on-farm collaborators Rufenacht Farms for the planting and harvesting of this plot. Thanks to summer agronomy intern Troy Grime for assistance with data collection. This project was supported by the Ohio Soybean Association Research and Education Fund.



THE OHIO STATE UNIVERSITY

For more information, contact:  
Eric Richer  
OSU Extension –Fulton County  
8770 State Route 108  
Wauseon, Ohio 43567  
[Richer.5@osu.edu](mailto:Richer.5@osu.edu)

